


Αναθέτουσα Αρχή:
**Υ.Π.Ε.ΧΩ.Δ.Ε. - Δ/νση Περιβαλλοντικού Σχεδιασμού Τμήμα
Διαχείρισης Φυσικού Περιβάλλοντος**

**Προσδιορισμός συμβατών δραστηριοτήτων
σε σχέση με τα είδη χαρακτηρισμού
των Ζωνών Ειδικής Προστασίας
της ορνιθοπανίδας**

Παραδοτέο 1 Κατάλογος βιβλιογραφικών πηγών



Ανάδοχος:
**Γραφείο Μελετών
ΤΑΣΟΣ ΔΗΜΑΛΕΞΗΣ**

Ιούλιος 2009

Ομάδα Μελέτης:

Τάσος Δημαλέξης	Δρ Βιολογίας	Επιστημονικός Υπεύθυνος του έργου
Θάνος Καστρίτης	Δρ Ωκεανογραφίας	Συντονισμός του έργου, Συντονισμός ομάδας οικολογικών απαιτήσεων
Κώστας Γρίβας	Γεωπόνος MSc	Συντονισμός ομάδας απειλών/μέτρων διατήρησης ειδών χαρακτηρισμού
Άρης Μανωλόπουλος	Περιβαλλοντολόγος	Σχεδιασμός και ανάπτυξη Βάσης Δεδομένων
Νίκη Καρδακάρη	Ορνιθολόγος	Ομάδα οικολογικών απαιτήσεων
Λευτέρης Κακαλής	Δασολόγος - Ορνιθολόγος	Ομάδα οικολογικών απαιτήσεων
Σταύρος Ξηρουχάκης	Δρ Βιολογίας	Ομάδα οικολογικών απαιτήσεων
Χρήστος Τσαϊτουρίδης	Δρ Νομικής	Νομικός σύμβουλος
Clairie Papazoglou	Regional Director of European Division, BirdLife International	Επιστημονικός σύμβουλος
Boris Barov	European Conservation Manager, BirdLife International	Επιστημονικός σύμβουλος

- Aarvak, T. & Øien, I. (2006). Δορυφορική παρακολούθηση της Νανόχηννας (*Anser erythropus*). *Οιωνός* 27: 28-33.
- Adamakopoulos, T., Gatzoyannis, S. & Poirazidis, C. (eds). (1995). *Special Environmental Study of Dadia Forest*. WWF Greece, Athens.
- Adamo, M.C., Puglisi, L. & Baldaccini, N.E. (2004). Factors affecting Bittern (*Botaurus stellaris*) distribution in a Mediterranean wetland. *Bird Conservation International* 14 (3): 153-164.
- Agostini, N., Cardelli, C. & Gustin, M. (2007). Factors shaping pathways of European Honey-buzzards (*Pernis apivorus*) during spring migration in the central Mediterranean basin. *Journal of Raptor Research* 41: 57-61.
- Alivizatos, H. & Bourdakis, S. (2002). Diet and Breeding Success of the Bonelli's Eagle (*Hieraetus fasciatus*) in Greece: Preliminary Data. *International Hawkwatcher* 5: 3-6.
- Alivizatos, H. & Goutner, V. (1997). Feeding habits of the Long-legged Buzzard (*Buteo rufinus*) during breeding in northeastern Greece. *Israel Journal of Zoology* 43: 257-266.
- Alivizatos, H., Goutner, V. & Karandinos, M.G. (1998). Reproduction and behaviour of the Long-legged Buzzard (*Buteo rufinus*) in north eastern Greece. *Vogelwarte* 39: 176-182.
- Alivizatos, H., Goutner, V. & Zogaris, S. (2005). Contribution to the diet of four owl species (Aves, Strigiformes), from mainland and island areas of Greece. *Belg. J. Zool.* 135(2): 109-118.
- Alivizatos, H., Papandropoulos, D. & Zogaris, S. (2004). Winter diet of the Greater Spotted Eagle (*Aquila clanga*) in the Amvrakikos wetlands, Greece. *Journal of Raptor Research* 38: 371-374.
- Amar, A. & Redpath, S.M. (2005). Habitat use by Hen Harriers *Circus cyaneus* on Orkney: Implications of land-use change for this declining population. *Ibis* 147: 37-47.
- Amcoff, M., Tjernberg, M. & Berg, A. (1994). Nest site choice of honey buzzard *Pernis apivorus*. *Ornis Svecica* 4: 145-158.
- Angelstam, P., Roberge, J-M, Lohmus, A., Bergmanis, M., Brazaitis, G., Donz-Breuss, M., Edenius, L., Kosinski, Z., Kurlavicius, P., Larmanis, V., Lukins, M., Mikusinski, G., Racinskis, E., Strazds, M. & Tryjanowski, P. (2004). Habitat modelling as a tool for landscape-scale conservation - a review of parameters for focal forest birds. In: Angelstam, P., Donz-Breuss, M. & Roberge, J-M (eds). Targets and tools for the maintenance of forest biodiversity. *Ecol. Bull.* 51: 427-453.
- Antoniou, V., Zantopoulos, N., Skartsi, T. & Tsoukali-Papadopoulou, H. (1996). Pesticide poisoning of animals of wild fauna. *Veterinary Human Toxicology* 38: 212-213.
- Arcos, J., Oro, D. & Sol, D. (2001). Competition between the yellow-legged gull *Larus cachinnans* and Audouin's gull *Larus audouinii* associated with commercial fishing vessels: The influence of season and fishing fleet. *Marine Biology* 139: 807-816.
- Arcos, J.M. & Oro, D. (1996). Changes in foraging range of Audouin's Gulls *Larus audouinii* in relation to a trawler moratorium in the western Mediterranean. *Waterbirds* 19: 128-131.
- Arroyo, B., Amar, A., Leckie, F., Buchanan, G.M., Wilson, J.D. & Redpath, S. (2009). Hunting habitat selection by hen harriers on moorland: Implications for conservation management. *Biological Conservation* 142: 586-596.
- Arroyo, B.E. & Bretagnolle, V. (1999). Breeding biology of the Short-eared Owl (*Asio flammeus*) in agricultural habitats of southwestern France. *Journal of Raptor Research* 33 (4): 287-294.
- Audsen, M. & Bateson, D. (2005). Winter cattle grazing to create foraging habitat for choughs *Pyrrhocorax pyrrhocorax* at South Stack RSBP Reserve, Anglesey, Wales. *Conservation Evidence* 2: 26-27.

- Aulen, G., & Lundberg, A. (1991). Sexual dimorphism and patterns of territory use by the Whitebacked Woodpecker *Dendrocopos leucotos*. *Ornis Scandinavica* 22: 60-64.
- Avilés, J.M. & Parejo, D. (2004). Farming practices and Roller *Coracias garrulus* conservation in south-west Spain. *Bird Conservation International* 14 (3): 173-181.
- Avilés, J.M., Sánchez, J.M. & Parejo, D.(2000). Nest-site selection and breeding success in the Roller (*Coracias garrulus*) in the Southwest of the Iberian peninsula. *Journal für Ornithologie* 141 (3): 345-350.
- Ayas, Z., Celikkan, H. & Aksu, M.L. (2008). Lead (Pb) and copper (Cu) concentration in the eggshells of Audouin's Gulls (*Larus audouinii*) in Turkey. *Turkish Journal of Zoology* 32: 379-384.
- Ακριώτης, Τ. (2008). *Α δημοσίευτα δεδομένα 1992-2008*.
- Ακριώτης, Τ., Ρήγας Ι. (1999). *Επιπτώσεις των βαρέων μετάλλων και των ιχνοστοιχείων στην Ορνιθοπανίδα*. Στο: Χατζηλάκου Δ. (Επιμέλεια). Συνοπτικός Οδηγός: Επιπτώσεις έργων και δραστηριοτήτων στα πουλιά και τους βιοτόπους τους. Διαχείριση βιοτόπων της ορνιθοπανίδας. Ελληνική Ορνιθολογική Εταιρεία, Αθήνα.
- Αλιβιζάτος, Χ. Π. (1996). *Βιολογία και οικολογία της Αετογερακίνας (Buteo rufinus CR.) στο Νομό Έβρου*. Διδακτορική διατριβή. Γεωπονικό Πανεπιστήμιο Αθηνών, Τμήμα Φυτικής Παραγωγής, Εργαστήριο Οικολογίας και Προστασίας Περιβάλλοντος. Αθήνα.
- Αλιβιζάτος, Χ., Ακριώτης, Τ., Χανδρινός, Γ. & Καζαντζίδης, Σ. (υπό έκδοση). *Οι Μεσοχειμωνιάτικες Καταμετρήσεις Υδροβίων Πουλιών στην Ελλάδα: 1967-2005*. Ελληνική Ορνιθολογική Εταιρεία & Ελληνικό Κέντρο Δακτυλίωσης Πουλιών.
- Αλιβιζάτος, Χ., Γκούτνερ, Β., Ρήγας, Ι., Αθανασιάδης, Α. & Ζόγκαρης, Σ. (2006). Χειμερινή οικολογία του Στικταετού (*Aquila clanga*) στο Δέλτα Έβρου και στους υγροτόπους Αμβρακικού. 3^ο Συνέδριο Ε.Οι.Ε & Ε.Ζ.Ε., Ιωάννινα. Σελ. 14-21.
- Bakaloudis D.E., Vlachos C.G. & Holloway G.J. (2005). Nest spacing and breeding performance in Short-toed Eagle *Circaetus gallicus* in northeast Greece. *British Trust for Ornithology, Bird Study* 52: 330-338.
- Bakaloudis, D.E., Vlachos C., Papageorgiou, N. & Holloway, G.J. (2001). Nest-site habitat selected by Short-toed Eagles *Circaetus gallicus* in Dadia forest (northeastern Greece). *Ibis* 143: 391-401.
- Bakaloudis, D.E., Vlachos C.G. & Holloway G.J. (1998). Habitat use by the Short-toed Eagle *Circaetus gallicus* and their reptilian prey during the breeding season in Dadia forest (North-Eastern Greece). *J. Ap. Ecology* 35: 821-828.
- Barbraud, C. & Mathevet, R. (2000). Is commercial reed harvesting compatible with breeding purple herons *Ardea purpurea* in the Camargue, southern France. *Environmental Conservation* 27 (4): 334-340.
- Barbraud, C., Lepley, M., Mathevet, R. & Mauchamp, A. (2002). Reedbed selection and colony size of breeding Purple Herons *Ardea purpurea* in southern France. *Ibis* 144 (2): 227-235.
- Barros, C. & De Juana, E. (1997). Breeding success of the Stone Curlew *Burhinus oedipnemos* at La Sercna (Badaioz, Spain). *Ardeola* 44 (2): 199-206.
- Baumann, S. (2000). Habitat structure and habitat use of the European Golden Orioles (*Oriolus oriolus*) during breeding and wintering. *Journal für Ornithologie* 141(2): 142-151.
- Bealey, C.E., Green, R.E., Robson, R., Taylor, C.R. & Winspear, R.(1999). Factors affecting the numbers and breeding success of Stone Curlews *Burhinus oedipnemos* at Porton Down, Wiltshire. *Bird Study* 46 (2): 145-156.
- Bechet, A., Ienmann, P. & Gaudin, R. (1998). Nest predation, temporal and spatial breeding strategy in the Woodchat Shrike *Lanius senator* in Mediterranean France. *Acta Oecologica* 19 (1): 81-87.

- Berg, A. (2008). Habitat selection and reproductive success of Ortolan Buntings *Emberiza hortulana* on farmland in central Sweden – the importance of habitat heterogeneity. *Ibis* 150: 565-573.
- BirdLife International. (2004). *Birds in Europe: population estimates, trends and conservation status*. BirdLife Conservation series No. 12. BirdLife International, Cambridge.
- BirdLife International. (2008). *Species factsheets*. Downloaded from <http://www.birdlife.org>.
- Blanco, G. (1997). Role of refuse as food for migrant, floater and breeding Black Kites (*Milvus migrans*). *Journal of Raptor Research* 31: 71-76.
- Blew, J., Hoffmann, M. & Nehls, G. (2007). Collision risks at sea: Species composition and altitude distributions of birds in Danish offshore wind farms. 2nd Scientific Conference on the Use of Offshore Wind Energy, Federal Ministry for the Environment, 20-21/2, Berlin.
- Bohus, M (2007). Breeding of the European roller (*Coracias garrulus*) in south-west Slovakia during 2001-2006. *Tichodroma* 19: 1-11.
- Bonetti, A. & Παπακωνσταντίνου, Κ. (2000). Τα πουλιά στη λιμνοθάλασσα Γιάλοβα και τη γύρω προστατευόμενη περιοχή. Στο: Καρδακάρη Ν. (ed.). Τα Πουλιά στο Διβάρι της Μεσσηνίας - Αναφορά στην ορνιθοπανίδα της περιοχής Natura 2000 Λιμνοθάλασσα της Πύλου και Νήσος Σφακτηρία. Ελληνική Ορνιθολογική Εταιρεία, Ευρωπαϊκή Επιτροπή, DGXI, Υ.ΠΕ.ΧΩ.Δ.Ε.
- Bonnaud, E., Bourgeois, K., Vidal, E., Legrand, J. & Le Corre M. (2009). How can the Yelkouan shearwater survive feral cat predation? A meta-population structure as a solution? *Population Ecology*. (in press).
- Bontzorlos, V.A., Peris, S.J., Vlachos, C.G. & Bakaloudis D.E. (2005). The diet of barn owl in the agricultural landscapes of central Greece. *Folia Zoologica* 54: 99-110.
- Bosch, J. (2003). Phenology and breeding parameters of Booted Eagle *Hieraaetus pennatus* in central Catalonia (Spain). *Ardeola* 50: 181-189.
- Bosch, J., Borrás, A. & Freixas J. (2005). Nesting habitat selection of Booted Eagle *Hieraaetus pennatus* in central Catalonia. *Ardeola* 52: 225-233.
- Bourdakis, S. & Xirouchakis S.M. (2009). The Bonelli's eagle (*Hieraaetus fasciatus*) in Greece. In: V. Hernandez (ed). The Bonelli's eagle. Ecology, behaviour and conservation. (in press).
- Bourdakis, S., Alivizatos, H., Azmanis, P., Hallmann, B., Panayotopoulou, M., Papakonstantinou, C., Probonas, N., Rousopoulos, Y., Skartsi, D., Stara, K., Tsiakiris, R. & Xirouchakis, S. (2004). The situation of Griffon Vulture in Greece. In: Slotta-Bachmayr, L., Bögel, R. & Camina, C.A (Eds.). The Eurasian Griffon Vulture (*Gyps fulvus*) in Europe and the Mediterranean. Status report and Action plan. EGVWG. Pp 48-56.
- Brambilla, M., Guidali, F. & Negri, I. (2008). The importance of an agricultural mosaic for Cirl Buntings *Emberiza cirlus* in Italy. *Ibis* 150: 628-632.
- Brambilla, M., Rubolini, D. & Guidali F. (2004). Rock climbing and Raven *Corvus corax* occurrence depress breeding success of cliff-nesting Peregrines *Falco peregrinus*. *Ardeola* 51: 425-430.
- Brambilla, M., Rubolini, D. & Guidali, F. (2007). Between land abandonment and agricultural intensification: Habitat preferences of Red-backed Shrikes *Lanius collurio* in low-intensity farming conditions. *Bird Study* 54 (2): 160-167.
- Brickle, N.W., Harper, D.G.C, Aebischer, N.J. & Cockayne, S.H. (2000). Effects of agricultural intensification on the breeding success of corn buntings *Miliaria calandra*. *Journal of Applied Ecology* 37: 742-755.
- Brooks, R. (1998). *Birding on the Greek Island of Lesbos*. Brookside Publishing. Fakenham, Norfolk.

- Brotons, L., Herrando, S., & Pons, P. (2008). Wildfires and the expansion of threatened farmland birds: the ortolan bunting *Emberiza hortulana* in Mediterranean landscapes. *Journal of Applied Ecology* 45: 1059-1066.
- Brotons, L., Wolff, A., Paulus, G. & Martin, J.M. (2005). Effect of adjacent agricultural habitat on the distribution of passerines in natural grasslands. *Biological Conservation* 124: 407-414.
- Brown, C. & Plug, I. (1990). Food choice and diet of the Bearded vulture *Gypaetus barbatus* in Southern Africa. *S. Afr. Zool.* 25 (3): 169-177.
- Brown, C.J. (1988). *A study of the Bearded vulture in Southern Africa*. Unpubl. Ph.D. Thesis. Univ. of Natal.
- Butler, R., Angelstam, P., Ekelund, P. & Schlaepfer, R. (2004). Dead wood threshold values for the three-toed woodpecker presence in boreal and sub-Alpine forest. *Biological Conservation* 119: 305-318.
- Βασιλειάδης, Β. προσωπική επικοινωνία
- Βλάχος, Χ. (1989). *Η οικολογία του Κραυγαετού (Aquila romana) στο Δάσος Δαδιάς του Νομού Έβρου*. Διδακτορική Διατριβή, Τμήμα Δασολογίας κα Φυσικού Περιβάλλοντος, ΑΠΘ. Σελ. 119.
- Callaghan, D.A. (1997). *European Union Species Action Plan: Ferruginous Duck (Aythya nyroca)*. Council of Europe Publication T-PVS/Birds (99) 2. Council of Europe Publishing, Strasbourg.
- Calvo, B. & Furness, R.W. (1995). Colony and nest-site Selection by Collared Pratincoles (*Glareola pratincola*) in Southwest Spain. *Colonial Waterbirds* 18 (1): 1-10.
- Camarda, A., Circella, E., Pennelli, D., Madio, A., Bruni, G., Lagrasta, V., Marzano, G., Mallia, E. & Campagnari E. (2006). Wild birds as biological indicators of environmental pollution: Biotyping and antimicrobial resistance patterns of *Escherichia coli* isolated from Audouin's gulls (*Larus audouinii*) living in the Bay of Gallipoli (Italy). *Italian Journal of Animal Science* 5: 287-290.
- Capizzi, D. (2000). Diet shifts of the tawny owl *Strix aluco* in central and northern Italy. *Italian Journal of Zoology* 67: 73-79.
- Carlson, A. (2000). The effect of habitat loss on a deciduous forest specialist species: the White-backed Woodpecker (*Dendrocopos leucotos*). *Forest Ecology and Management* 131: 215-221.
- Carrete, M., Sanchez-Zapata, J.A., Martinez, J.E. & Calvo, J.E. (2002). Predicting the implications of conservation management: A territorial occupancy model of Bonelli's eagle in Murcia, Spain. *Oryx* 36: 349-356.
- Carss, D.N. (1993). Shags *Phalacrocorax aristotelis* at cage fish farms in Argyll, western Scotland. *Bird Study* 40: 203-211.
- Carss, D.N. (1994). Killing of piscivorous birds at Scottish fin fish farms, 1984-87. *Biological Conservation* 68: 181-188.
- Casado, E. & Ferrer M. (2005). Analysis of reservoir selection by wintering Ospreys (*Pandion haliaetus*) in Andalusia, Spain: A potential tool for reintroduction. *Journal of Raptor Research* 39: 168-173.
- Castellanos-Vera, A. & Rivera, E. (2007). Hunting patterns and success of an osprey (*Pandion haliaetus*) population at Magdalena Bay, Baja California Sur, Mexico. *Ciencias Marinas* 33: 325-333.
- Catry, T., Ramos, J.A., Catry, I., Allen-Revez, M. & Grade, N. (2004). Are salinas a suitable alternative breeding habitat for Little Terns *Sterna albifrons*? *Ibis* 146 (2): 247-257.
- Catsadorakis, G. & Källander, H. (1999). Densities, habitat and breeding parameters of the Sombre Tit *Parus lugubris* in Prespa National Park, Greece. *Bird Study* 46: 373-375.

- Ceballos, O. & Donazar, J.A. (1989). Factors influencing the breeding density and nest-site selection of the Egyptian Vulture (*Neophron percnopterus*). *J. Ornithology* 130: 353-359.
- Chamberlain, D.E. & Crick, H.Q.P. (1999). Population declines and reproductive performance in skylarks *Alauda arvensis* in different regions and habitats of the United Kingdom. *Ibis* 141: 38-51.
- Chamberlain, D.E., Wilson, A.M., Browne, S.J. & Vickery, J.A. (1999). Effects of habitat type and management on the abundance of skylarks in the breeding season. *Journal of Applied Ecology* 36: 856-870.
- Charter, M., Izhaki, I., Shapira, L. & Leshem Y. (2007). Diets of urban breeding barn owls (*Tyto alba*) in Tel Aviv, Israel. *Wilson Journal of Ornithology* 119: 484-485.
- Cormier, J-P., Fustec, J., Pithon, J. & Choisy P. (2008). Selection of nesting habitat by Montagu's Harriers *Circus pygargus* and Hen Harriers *Circus cyaneus* in managed heaths. *Bird Study* 55: 86-93.
- Cramp, S. & Perrins, C.M. (eds). (1993). *Handbook of the birds of Europe, the Middle East and Africa: The birds of the Western Palearctic*. Oxford University Press.
- Crivelli, A. J., Catsadorakis G., Hatzilacou D. & Naziridis T. (1997 a). *Pelecanus crispus* Dalmatian Pelican. BWP (Birds of Western Palearctic) Update 1: 149-153.
- Crivelli, A. J., Catsadorakis, G. & Naziridis T. (1997 b). *Pelecanus onocrotalus* Great White Pelican. BWP (Birds of Western Palearctic) Update 1: 144-148.
- Crivelli, A., Μαλακού Μ., Χατζηλάκου, Δ. & Ναζηριδης, Θ. Προσωπική επικοινωνία.
- Croxall, J.P., Evans, P.G.H. & Schreiber R.W. (1984). *Status and conservation of the world's seabirds*. ICPB, Technical Pub. 2., Cambridge.
- Czeszczewik, D. & Walankiewicz, W. (2006). Logging affects the white-backed woodpecker *Dendrocopos leucotos* distribution in the Białowieza Forest. *Annales Zoologici Fennici* 43: 221-227.
- Dale, S. & Olsen, B.F.G. (2002). Use of farmland by ortolan buntings (*Emberiza hortulana*) nesting on a burned forest area. *Journal für Ornithologie* 143: 133-144.
- De La Montaña, E., Rey-Benayas, J.M. & Carrascal, L.M. (2006). Response of bird communities to silvicultural thinning of Mediterranean maquis. *Journal of Applied Ecology* 43: 651-659.
- De Leon, A. & Minguez, E. (2003). Occupancy rates and nesting success of European Storm-petrels breeding inside artificial nest-boxes. *Scientia Marina* 67: 109-112. Suppl. 2.
- Denker, E., Buthe, A., Glimm, D., Holker, M., Prunte, W. & Trendelkamp, T. (2003). Changes in the DDT and PCB burden in life stages of Montagu's (*Circus pygargus*) and Marsh Harriers (*Circus aeruginosus*) from North-Rhine Westfalia, Germany. *Journal für Ornithologie* 144: 411-417.
- Dennis, T.E. (2007). Reproductive activity in the Osprey (*Pandion haliaetus*) on Kangaroo Island, South Australia. *Emu* 107: 300-307.
- Digby, R. & Milwright, P. (1998). Breeding biology of the Golden Oriole *Oriolus oriolus* in the fenland basin of the Eastern Britain. *Bird Study* 45(3): 320-330.
- Dimalexis, A., Xirouchakis, S., Latsoudis, P., Portolou, D., Karris, G., Georgiakakis, P., Fric, J., Barboutis, C., Bourdakis, S., Kakalis, E., Ivovič, M. & Kominos, T. (2007). Breeding distribution and population status of the Eleonora's falcon (*Falco eleonora*) in Greece. *Journal of Ornithology* 149: 23-30.
- Donald, P.F., Wilson, J.D. & Shepherd, M. (1994). The decline of the corn bunting. *British Birds* 87: 106-132.
- Donazar, J.A. (1993). *Los Buitres Ibericos, Biología y Conservacion*. Reyero, J.M. (ed.) Madrid.

- Δρετάκης, Μ. & Παπακωνσταντίνου, Κ. (2008). Οι γλάροι στην Ελλάδα. Οικωνός 34: 22-26.
- European Commission, (2007). *Management Plan for Curlew (Numenius arquata)*. Luxembourg: Office for Official Publications of the European Communities.
- European Union Management Plan for Golden Plover *Pluvialis apricaria* (2008-2010).
- European Union Management Plan for Lapwing *Vanellus vanellus* (2008-2010).
- Evans, A.D. & Smith, K.W. (1994). Habitat selection of circl buntings *Emberiza circlus* wintering in Britain. *Bird Study* 47: 81-87.
- Evans, A.D., Smith, K.W., Buckingham, D.L. & Evans, J. (1997). Seasonal performance and nestling diet of Circl Buntings *Emberiza circlus* in England. *Bird Study* 44: 66-79.
- Ελληνική Ορνιθολογική Εταιρεία, βάση δεδομένων.
- Επιτροπή Αξιολόγησης Ορνιθολογικών Παρατηρήσεων (ΕΑΟΠ), βάση δεδομένων.
- Fasola, M. (1993). *Distribution, population and Habitat Requirements of the Common Tern and the Little Tern breeding in the Mediterranean*. In: Aguilar, J.S., Monbailliu, X. & Paterson, A.M. Status and Conservation of Seabirds, Proceedings of the 2nd Medmaravis, SEO, Madrid.
- Fasola, M., Goutner, V. & Walmsley, J. (1993). *Comparative breeding Biology of the Gulls and Terns in the four main Deltas of the Northern Mediterranean*. In: Aguilar, J.S., Monbailliu, X. & Paterson, A.M. Status and Conservation of Seabirds. Proceedings of the 2nd Medmaravis, SEO, Madrid.
- Fayt, P. (1999). Available insect prey in bark patches selected by the Three-toed Woodpecker *Picoides tridactylus* prior to reproduction. *Ornis Fennica* 76: 135-140.
- Ferguson-Lees, J. & Christie, D.A. (2001). *Raptors of the world*. Helm, London.
- Fernandez, C. & Azkona, P. (1996). Influence of forest structure on the density and distribution of the White-backed Woodpecker *Dendrocopos leucotos* and black Woodpecker *Dryocopus martius* in Quinto Real (Spanish western Pyrenees). *Bird Study* 43: 305-313.
- Fonderflick, J., Thevenot, M. & Guillaume, C.P. (2005). Habitat of the ortolan bunting *Emberiza hortulana* on a cause in southern France. *Vie et Milieu – Life and Environment* 55: 109-120.
- Fox, A.D. & Heldbjerg, H. (2008). Which regional features of Danish agriculture favour the corn bunting in the contemporary farming landscape? *Agriculture, Ecosystems and Environment* 126: 261-269.
- Fox, T.(A.D.), Desholm, M., Kahlert, J., Petersen, K. & Christensen, T.K. (2007). *Avian avoidance behaviour and collision risk: Results from post construction monitoring of the Danish offshore wind farms at Horns Rev and Nysted*. 2nd Scientific Conference on the Use of Offshore Wind Energy, Federal Ministry for the Environment, 20-21/2, Berlin.
- Francour, P. & Thibault, J.C. (1996). The diet of breeding Osprey *Pandion haliaetus* on Corsica: Exploitation of a coastal marine environment. *Bird Study* 43: 129-133.
- Fuller, R.J., Gregory, R.D., Gibbons, D.W., Marchant, J.H., Wilson, J.D., Baillie, S.R. & Carter, N. (1995). Population declines and range contractions among lowland farmland birds in Britain. *Conservation Biology* 9: 1425-1441.
- Gallo-Orsi, U. (2003). Species Action Plans for the conservation of seabirds in the Mediterranean Sea: Audouin's gull, Balearic shearwater and Mediterranean shag. *Scientia Marina* 67 (Supplement 2): 47-55.
- Garcia-Dios, I.S. (2006). Diet of the booted eagle *Hieraaetus pennatus* in the south of Avila: Importance of the songbirds. *Ardeola* 53: 39-54.
- Garmendia, A., Carcamo, S. & Schwendtner, O. (2006). Forest management considerations for conservation of Black Woodpecker *Dryocopus martius* and White-backed

- Woodpecker *Dendrocopos leucotos* populations in Quinto Real (Spanish Western Pyrenees). *Biodiversity and Conservation* 15: 1399-1415.
- Garthe, S. & Huppopp, O. (2004). Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. *Journal of applied Ecology* 41(4): 724-734.
- Gensbol, B. & Thiede, W. (2008). *Birds of Prey*. Collins.
- Giannangeli, L., De Sanctis, A., Manginelli, R. & Medina, F.M. (2005). Seasonal variation of the diet of the stone curlew *Burhinus oedicnemus distinctus* at the Island of La Palma, Canary Islands. *Ardea* 92(2): 175-184.
- Gjerde, I., Saetersdal, M. & Nilsen, T. (2005). Abundance of two threatened woodpecker species in relation to the proportion of spruce plantations in native pine forests of western Norway. *Biodiversity and Conservation* 14: 377-393.
- Golawski, A. & Dombrowski, A. (2002). Habitat use of Yellowhammers *Emberiza citrinella*, Ortolan Buntings *Emberiza hortulana* and Corn Buntings *Miliaria calandra* in farmland of east-central Poland. *Ornis Fennica* 79: 164-172.
- Goławski, A. & Goławska, S. (2008). Habitat preference in territories of the red-backed shrike *Lanius collurio* and their food richness in an extensive agriculture landscape. *Acta Zoologica Academiae Scientiarum Hungaricae* 54(1): 89-97.
- Golawski, A. & Meissner, W. (2008). The influence of territory characteristics and food supply on the breeding performance of the Red-Backed Shrike (*Lanius collurio*) in an extensively farmed region of eastern Poland. *Ecol. Res.* 23: 347-353.
- Goławski, A. (2008). No evidence of weather effect found on the clutch size, eggs sizes and their hatchability in the red-backed shrike *Lanius collurio* in eastern Poland. *Annales Zoologici Fennici* 45(6): 513-520.
- González, L.M., Margalida, A., Sánchez, R. & Oria J. (2006). Supplementary feeding as an effective tool for improving breeding success in the Spanish imperial eagle (*Aquila adalberti*). *Biological Conservation* 129: 477-486.
- Goodman, S.M. & Haynes, C.V. (1992). The diet of the lanner (*Falco biarmicus*) in a hyper-arid region of the eastern Sahara. *Journal of Arid Environments* 22: 93-98.
- Goutner, V. & Handrinos, G. (1990). The occurrence of Slender-billed Curlews (*Numenius tenuirostris*) in Greece. *Biol. Conserv.* 53: 47-60.
- Goutner, V. (1987). Vegetation preferences by colonies of Mediterranean Gulls (*Larus melanocephalus*) and Gull-billed Terns (*Gelochelidon nilotica*) in the Evros Delta. *Seevogel* 8: 29-31.
- Goutner, V. (1997). Use of the Drana Lagoon (Evros Delta, Greece) by Threatened Colonially Nesting Waterbirds and its Possible Restoration. *Biological Conservation* 81: 113-120.
- Goutner, V., Albanis, T. & Konstantinou, L. (2005). PCBs and organochlorine pesticide residues in eggs of threatened colonial charadriiform species (Aves, Charadriiformes) from wetlands of international importance in northeastern Greece. *Belgian Journal of Zoology* 135(2): 157-163.
- Goutner, V., Charalambidou, T., & Albanis, A. (1997). Organochlorine Insecticide Residues in Eggs of the Little Tern (*Sterna albifrons*) in the Axios Delta, Greece. *Bull. Environmental Contamination and Toxicology* 58: 61-66.
- Goutner, V., Portolou, D., Papakonstantinou, K., Tsiakiris, R., Pavlidis, A., Zogaris, S., Kominos, T., Galanaki, A. & Oro, D. (2000). Nest site characteristics of Audouin's Gull in the eastern Mediterranean. *Waterbirds* 23: 74-83.
- Green, R.E., Tyler, G.A. & Bowden, C.G.R. (2000). Habitat selection, ranging behaviour and diet of the stone curlew (*Burhinus oedicnemus*) in southern England. *Journal of Zoology* 250(2): 161-183.

- Gremillet, D., Argentin, G., Schulte, B. & Culik B.M. (1998). Flexible foraging techniques in breeding Cormorants *Phalacrocorax carbo* and Shags *Phalacrocorax aristotelis*: Benthic or pelagic feeding? *Ibis* 140: 113-119.
- Gretton, A. (1991). *Conservation of the Slender-billed Curlew*. ICBP Monograph no 6. Cambridge. Pp 159.
- Grubač, B. (1985). The death of a female Bearded Vulture (*Gypaetus barbatus*) in Macedonia. *Bulletin "Gypaetus barbatus"* 7: 26-27.
- Grzybek, J., Michalak, I., Osiejuk, T.S. & Tryjanowski, P. (2008) Densities and habitats of the Tawny Pipit *Anthus campestris* in the Wielkopolska region (W. Poland). *Acta Ornithologica* 43(2): 221-225.
- Guerrieri, G., Pietrelli, L. & Biondi, M. (1996). Status and reproductive habitat selection of three species of Shrikes, *Lanius collurio*, *L. senator* and *L. minor* in a Mediterranean area. (Proc. of the First Intern. Shrike Symposium Found). *Vert. Zool.* 6: 167-171.
- Hakkinen, H. & Hasanen, E. (1980). Mercury in eggs and nestlings of the osprey (*Pandion haliaetus*) in Finland and it's bioaccumulation from fish. *Annales Zoologici Fennici* 17: 131-139.
- Hald, A.B. (1999). The impact of changing the season in which cereals are sown on the diversity of the weed flora in rotational fields in Denmark. *Journal of Applied Ecology* 36: 24-32.
- Hallmann B. (1989). Status and distribution of the Aquila in Greece. *Biol. Gallo-hellenica* 15: 171-176.
- Hallmann, B. (1996). *Greece's Endangered Birds of Prey, Eleven hour to 10 species*. WWF - Hellas Technical report. Athens.
- Hallmann, B. (1996). *The decline of the Imperial Eagle Aquila heliaca in Greece*. In: Eagle Studies, Meyburg, B-U & R.D. Chancellor, World Working Group for Birds of Prey and Owls. Berlin, Pp 439-442.
- Hancock, J. & Kushlan, J. (1984). *The Herons Handbook*. CroomHelm, London.
- Handrinos, G. & Goutner, V. (1990). On the occurrence of the Lesser White-fronted Goose (*Anser erythropus*) in Greece. *J. Orn.* 131: 160-165.
- Handrinos, G. (1985). *The status of vultures in Greece*. In: Conservation Studies of Raptors. Newton, I. & Chancellor, R. (Eds.): 103-115. ICBP Technical Publication No 5. ICBP. Cambridge.
- Handrinos, G. (1987). *The Golden Eagle in Greece*. Actes 1er Coll. Intern. Aigle Royal en Europe, Arvieux, June 1986, Pp 18-22
- Handrinos, G. (1991). *The status of geese in Greece*. In: Fox, A.D., Madsen, J. & Van Rhijn J. (eds.) Western Palearctic Geese. Proc. IWRB Symposium, Kleve, 1989. *Ardea* 79(2): 175-178.
- Handrinos, G. (1995). The White-headed Duck (*Oxyura leucocephala*) in Greece. IWRB Threatened Waterfowl. *Res. Group. Newsl.* 7: 6-7.
- Handrinos, G., & Akriotis, T. (1997). *The birds of Greece*. C. Helm, A & C Black, London.
- Heath, S.R., Servello, F.A. (2008). Effects of predation and food provisioning on Black Tern chick survival. *Wilson Journal of Ornithology* 120(1): 167-175.
- Hegemann, A. (2006). Successful breeding of a Common Buzzard *Buteo buteo* in a cliff nest inside a quarry in Northwest Germany. *Vogelwarte* 44: 131-133.
- Heneberg, P. & Simecek, K. (2004). Nesting of European bee-eaters (*Merops apiaster*) in Central Europe depends on the soil characteristics of nest sites. *Biologia - Section Zoology* 59(2): 205-211.
- Heneberg, P. (2008). Soil penetrability as a key factor affecting the nesting of burrowing birds. *Ecological Research*: 1-7.

- Heredia, B. & Heredia, R. (1997). *European Union Species Action Plan for the Lammergeyer (Gypaetus barbatus)*. European Union Action Plans for 8 Priority Birds Species.
- Heredia, R. & B. (Eds.). (1991). *El Quebrantahuesos (Gypaetus barbatus) en los Pirineos. Características ecológicas y biología de la conservación*. ICONA, Madrid.
- Hernandez, M. & Margalida, A. (2008). Poison - related mortality effects in the endangered Egyptian vulture (*Neophron percnopterus*) population in Spain. *Eur. J. Wildlife Research* DOI 10.1007/s10344-009-0255-6.
- Hoi, H., Hoi, C., Kristofik, J. & Darolova, A. (2002). Reproductive success decreases with colony size in the European bee-eater. *Ethology, Ecology and Evolution* 14 (2): 99-110.
- Hölzinger, J. (1987). *Bird of Baden -Württemberg, 1: threat and conservation*. Ulmer Verlag, Karlsruhe.
- Houston, D.C. (1976). Breeding of the white - backed and Rueppel's Griffon vultures, *Gyps africanus* and *G. rueppellii*. *Ibis* 118: 14-40.
- Hughes, B., Robinson, J.A., Green, A.J., Li, Z.W.D. & Mundkur, T. (Compilers). 2006. International Single Species Action Plan for the Conservation of the White-headed Duck *Oxyura leucocephala*. CMS Technical Series No. 13 & AEWA Technical Series No.8. Bonn, Germany.
- Imbeau, L., & Desrochers, A. (2002). Area sensitivity and edge avoidance: the case of the Three-toed Woodpecker (*Picoides tridactylus*) in a managed forest. *Forest Ecology and Management* 164: 249-256.
- Imbeau, L., Monkkonen, M. & Desrochers, A. (2001). Long-term effects of forestry on birds of the eastern Canadian boreal forests: a comparison with Fennoscandia. *Conserv. Biol.* 15: 1151-1162.
- Íñigo, A., Infante, O., Valls, J. & Atienza, J.C. 2008. *Directrices para la redaccion de planes o instrumentos de gestion de las Zonas de Especial Proteccion para las Aves*. SEO/ Birdlife, Madrid.
- Isenmann, P. & Debout, G. (2000). Vineyards harbour a relict population of Lesser Grey Shrike (*Lanius minor*) in Mediterranean France. *Journal für Ornithologie* 141 (4): 435-440.
- Isenmann, P. & Fradet, G. (1995). Is the nesting association between the Orphean Warbler (*Sylvia hortensis*) and the Woodchat Shrike (*Lanius senator*) an anti-predator oriented mutualism? *J. Orn.* 136: 288-291.
- Isenmann, P. & Goutner, V. (1993). *Breeding Status of the Slender-billed Gull (Larus genei) in the Mediterranean Basin*. In: Aguilar, J.S., Monbailliu, X. & Paterson, A.M. Status and Conservation of Seabirds, Proceedings of the 2nd Medmaravis, SEO, Madrid.
- Jenkins, A.R. & Avery, G.M. (1999). Diets of breeding Peregrine and Lanner Falcons in South Africa. *J. Rap.Research* 33: 190-206.
- Jerrentrup, H. & Χανδρινός Γ. (1987). Η βαρυχειμωιά του 1984/85 και οι επιπτώσεις της στα υδρόβια πουλιά. *Η Φύση* 36: 9-12.
- Jerrentrup, H. (1988). *White-tailed Eagle: Population developments and threats in the eastern Mediterranean. Proposals for conservation in Greece*. In: Ornis Consult (ed). Conservation and Management of the White-tailed Eagle *Haliaeetus albicilla* in the European Community, Copenhagen.
- Johnson, D.H. & Igl, L.D. (2001). Area requirements of grassland birds: A regional perspective. *Auk* 118: 24-34.
- Johnson, M.D. & Horn, C.M. (2008). Effects of rotational grazing on rodents and raptors in a coastal grassland. *Western North American Naturalist* 68 (4): 444-452.
- Jones, T., Martin, K., Barov, B., Nagy, S. (Compilers). 2008. International Single Species Action Plan for the Conservation of the Western Palearctic Population of the Lesser White-fronted Goose *Anser erythropus*. AEWA Technical Series No.36. Bonn, Germany.

- Joris, E. & Stienen, E. (2009). *Impact of wind Turbines on Terns in Zeebrugge, Belgium*. In: Stienen Eric, Norman Ratcliffe, Jan Seys, Jurgen Tack, Jan Mees and Ingrid Dobbelaere (eds.). 2009. Seabird Group 10th International Conference. Provincial Court, Brugge, 27-30 March 2009. VLIZ Special Publication 42. Communications of the Research Institute for Nature and Forest- INBO.M.2009.1. Research Institute for Nature and Forest (INBO), Brussels, Belgium-Flanders Marine institute (VLIZ). Oostende, Belgium. Viii+68 p.
- Kaisanlahti-Jokimäki, M-L, Jokimäki, J., Huhta, E., Ukkola, M., Helle, P. & Ollila, T. (2008). Territory occupancy and breeding success of the Golden Eagle (*Aquila chrysaetos*) around tourist destinations in northern Finland. *Ornis Fennica* 85: 2-12.
- Kakalis, E. & Akriotis, T. (2007). *Nest site characteristics of Krüper's Nuthatch Sitta krueperi on the island of Lesbos, Greece*. 2nd International Congress of Eurasian Ornithology, Antalya, Turkey, 26-29 Oct 07 (poster presentation).
- Karauz, S., Kiraç, C.O. & Eken, G. (2000). *Mediterranean Gull, Slender-billed Gull and Gull-billed Tern in Turkey*. In: Yésou, P. & Sultana, J. Monitoring and Conservation of Birds, Mammals and Sea Turtles of the Mediterranean and Black Seas. Proceedings 5th Medmaravis Symposium. Environment Protection Dept., Malta.
- Katzner, T.E., Bragin, E.A., Knick, S.T. & Smith, A.T. (2006). Spatial structure in the diet of imperial eagles *Aquila heliaca* in Kazakhstan. *Journal of Avian Biology* 37: 594-600.
- Kazantzidis, S. & Goutner, V. (2008). Abundance and habitat use by herons (Ardeidae) in the Axios Delta, northern Greece. *Journal of Biological Research-Thessaloniki* 10: 129-138.
- Kazantzidis, S. & Koutrakis, E. (2007). Great Cormorants and fisheries in Greece: conflict & management. NAGREF. Unp. Report. Pp 4.
- Kosinski, Z. & Winiiecki, A. (2004). Nest-site selection and niche partitioning among the great spotted woodpecker *Dendrocopos major* and middle spotted woodpecker *Dendrodopos medius* in riverine forest of central.Europe. *Ornis Fenn.* 81: 145-156.
- Kosinski, Z. & Winiiecki, A. (2005). Factors affecting the density of the middle spotted woodpecker *Dendrocopos medius*: a macrohabitat approach. *J. Ornithol.* 146: 263-270.
- Kristin, A., Hoi, H., Valera, F. & Hoi, C. (2007). Philopatry, dispersal patterns and nest-site reuse in Lesser Grey Shrikes (*Lanius minor*). *Biodivers. Conserv.* 16: 987-995.
- Kristin, A., Hoi, H., Valera, F. & Hoi, C. (2007). The importance of breeding density and breeding synchrony for paternity assurance strategies in the lesser grey shrike. *Folia Zoologica* 57(3): 240-250.
- Kristin, A., Hoi, H., Valera, F. & Hoi, H. (2000). Breeding biology and breeding success of the Lesser Grey Shrike (*Lanius minor*) in a stable and dense population. *Ibis* 142(2): 305-311.
- Krüger, O. (2002). Analysis of nest occupancy and nest reproduction in two sympatric raptors: Common buzzard *Buteo buteo* and goshawk *Accipiter gentiles*. *Ecography* 25: 523-532.
- Καζαντζίδης, Σ. & Νοϊδου, Μ. (συντον. έκδοσης). (2008). Προσδιορισμός της Φαινολογίας Μετανάστευσης των Θηρευσίμων Υδρόβιων Πουλιών. ΥΠ.Α.Α.Τ - Γεν. Δ/ση Δασών, ΕΘΙΑΓΕ/ Ι.Δ.Ε. Θεσσαλονίκη. 255 σ.+Παράρτημα.
- Καζαντζίδης, Σ. (2005). *Ερωδιοί*. Ελληνική Ορνιθολογική Εταιρεία
- Καζαντζίδης, Σ. προσωπική επικοινωνία.
- Κακαλής, Ε. (2004). Κατανομή, πληθυσμιακή πυκνότητα και επιλογή ενδιαιτήματος του Τουρκοτσοπανάκου (*Sitta krueperi*) στη Λέσβο. Μεταπτυχιακή εργασία. Τμήμα Περιβάλλοντος, Πανεπιστήμιο Αιγαίου.
- Κακαλής, Ε., αδημοσίευτα στοιχεία.
- Καρδακάρη, Ν., Λατσούδης, Π. & Παπανδρόπουλος, Δ. (2006). Προστασία φωλεοποίησης στα υλίπεδα του Μετοχίου: Κατασκευή και τοποθέτηση ηλεκτρικής περίφραξης. Σελ 47-51.

- Στο: Έκθεση δικτύωσης - Διαχείριση Νερού, Χλωρίδας και Πανίδας στους Ελληνικούς Υγρότοπους: Εμπειρίες από τα έργα LIFE-ΦΥΣΗ ΙΙΙ. Νομαρχιακή Αυτοδιοίκηση Δράμας - Καβάλας - Ξάνθης. Ξάνθη. Σελ. 64.
- Κατσαδωράκης, Γ. & Παραγκαμιάν, Κ. (2007). Απογραφή των υγροτόπων νησιών του Αιγαίου: Ταυτότητα, οικολογική κατάσταση και απειλές. Παγκόσμιο Ταμείο για τη Φύση - WWF Ελλάς, Αθήνα. Σελ. 392.
- Κολιόπουλος, Γ. (1999). Επιπτώσεις των παρασιτοκτόνων στην Ορνιθοπανίδα. Στο: Χατζηλάκου Δ. (Επιμέλεια). Συνοπτικός Οδηγός. Επιπτώσεις έργων και δραστηριοτήτων στα πουλιά και τους βιοτόπους τους. Διαχείριση βιοτόπων της ορνιθοπανίδας. Ελληνική Ορνιθολογική Εταιρεία, Αθήνα.
- Κουτσερή, Ε., προσ. επικοινων.
- Leckie, F.M., Arroyo, B.E., Thirgood, S.J. & Redpath S.M. (2008). Parental differences in brood provisioning by Hen Harriers *Circus cyaneus*. *Bird Study* 55: 209-215.
- Lepley, M., Ranc, S., Isenmann, P., Bara, T., Ponel, P. & Guillemain, M. (2004). Diet and gregarious breeding in lesser Grey Shrike (*Lanius minor*) in Mediterranean France. *Revue d'Ecologie (La Terre et la Vie)* 59(4): 591-602.
- Lilliendahl, K. & Solmundsson, J. (2006). Feeding ecology of sympatric European shags *Phalacrocorax aristotelis* and great cormorants *P. carbo* in Iceland. *Marine Biology* 149: 979-990.
- Liven-Schulman, I., Leshem, Y., Alon, D. & Yom-Tov, Y. (2004). Causes of population declines of the Lesser Kestrel *Falco naumanni* in Israel. *Ibis* 146: 145-152.
- Lohmus, A. (2005) Are timber harvesting and conservation of nest sites of forest-dwelling raptors always mutually exclusive? *Animal Conservation* 8: 443-450.
- López-López, P., García-Ripollés, C. & Urios, V. (2007). Population size, breeding performance and territory quality of Bonelli's Eagle *Hieraetus fasciatus* in eastern Spain. *Bird Study* 54: 335-342.
- Mahmood-ul-Hassan, M., Beg, M.A. & Mushtaq-ul-Hassan, M. (2007). Locality related changes in the diet of the Barn Owl (*Tyto alba stertens*) in agroecosystems in central Punjab, Pakistan. *Ison Journal of Ornithology* 119: 479-483.
- Mahmood-ul-Hassan, M., Beg, M.A., Mushtaq-ul-Hassan, M., Mirza, H.A. & Siddique, M. (2007). Nesting and diet of the Barn Owl (*Tyto alba*) in Pakistan. *Journal of Raptor Research* 41: 122-129.
- Manganaro, A., Pucci, L., Ranazzi, L. & Salvati, L. (2001). Frogs as prey of tawny owls *Strix aluco* in Central Italy: A comparison with diets from Central Europe. *Alauda* 69: 550-552.
- Manosa, S., Oro, D. & Ruiz, X. (2004). Activity patterns and foraging behaviour of Audouin's gulls in the Ebro Delta, NW Mediterranean. *Scientia Marina* 68: 605-614.
- Marniche, F., Voisin, J-F., Doumandji, S. & Baziz, B. (2007). Diet of the European Bee-eater *Merops apiaster* in the Ichkeul National Park (Tunisia). *Revue d'Ecologie (La Terre et la Vie)* 62(4): 351-362.
- Marquiss, M., Robinson, L. & Tindal, E. (2007). Marine foraging by Ospreys in southwest Scotland: Implications for the species' distribution in Western Europe. *British Birds* 100: 456-465.
- Martinez, J.E., Pagan, I. & Calvo J.F. (2006). Factors influencing territorial occupancy and reproductive output in the Booted Eagle *Hieraetus pennatus*. *Ibis* 148: 807-819.
- Martinez-Lopez, E., Maria-Mofjica, P., Martinez, J.E., Calvo, J.F., Wright, J., Shore, R.F., Romero, D. & Garcia-Fernandez, A.J. (2007). Organochlorine residues in booted eagle (*Hieraetus pennatus*) and goshawk (*Accipiter gentilis*) eggs from southeastern Spain. *Environmental Toxicology and Chemistry* 26: 2373-2378.

- Martinez-Lopez, E., Romero, D., Maria-Mojica, P., Martínez, J.E., Calvo, J.F. & Garcia-Fernandez, A.J. (2009). Changes in blood pesticide levels in booted eagle (*Hieraaetus pennatus*) associated with agricultural land practices. *Ecotoxicology and Environmental Safety* 72: 45-50.
- Masero, J.A., Perez-Hurtado, A., Castro, M. & Arroyo, G.M. (2000). Complementary use of intertidal mudflats and adjacent salinas by foraging waders. *Ardea* 88(2): 177-191.
- Mason, C.F. & MacDonald, S.M. (2000). Corn bunting *Miliaria calandra* populations, landscape and land-use in an arable district of eastern England. *Bird Conserv. Int.* 10: 169-186.
- Massero, J.A. (2003). Assessing alternative anthropogenic habitats for conserving waterbirds: salinas as buffer areas against the impact of natural habitat loss for shorebirds. *Biodiversity and Conservation* 12(6): 1157-1173.
- McConnell, S., O'Connell, T.J. & D.M. & Leslie, J.R. (2008). Land cover associations of nesting territories of three sympatric buteos in shortgrass prairie. *Wilson Journal of Ornithology* 120: 708-716.
- Medeiros, R., Ramos, J., Paiva, V., Almeida, A., Pedro, P. & Antunes, S. (2007). Signage reduces the impact of human disturbance on little tern nesting success in Portugal. *Biological Conservation* 135: 99-100.
- Meunier, F.D., Verheyden, C. & Jouventin P. (2000). Use of roadsides by diurnal raptors in agricultural landscapes. *Biological Conservation* 92: 291-298.
- Michelat, D. & Giraudoux, P.(2000). The feeding behaviour of breeding Short-Eared Owls (*Asio flammeus*) and relationships with communities of small mammal prey. *Revue d'Ecologie (La Terre et la Vie)* 55(1): 77-91.
- Mikkola, H. (1983). *The Owls of Europe*. T. & A.D. Poyser, London.
- Monbailliu, X. (2009). *Bird Species to be added to Annex II of the Specially Protected Areas, Barcelona Convention proposed amendments to the Specially Protected Areas and biological diversity protocol*. United Nations Environment Programme, Mediterranean Action-Plan, Regional Activity Centre for Specially Protected Areas.
- Monson, C.S. (2001). Ground-nesting ospreys in Utah. *Journal of Raptor Research* 35: 257-258.
- Morris, A. J., Holland, J.M., Smith, B. & Jones, N. E. (2004). Sustainable Arable Farming For an Improved Environment (SAFFIE): managing winter wheat sward structure for Skylarks *Alauda arvensis*. *Ibis* 146 (Suppl. 2): 155-162.
- Moskat, C. & Fuisz, T.I. (2002). Habitat segregation among the woodchat shrike *Lanius senator*, the red-backed shrike, *Lanius collurio* and the masked shrike, *Lanius nubicus*, in NE Greece. *Folia Zool.* 51(2): 103-111.
- Movalli, P.A. (2000). Heavy metal and other residues in feathers of laggar falcon *Falco biarmicus* jugger from six districts of Pakistan. *Env. Pollution* 109: 267-275.
- Mullarney, K., Svensson, L., Zetterstrom, D. & Grant, P. (1999). Τα Πουλιά της Ελλάδας, της Κύπρου και της Ευρώπης. Ελληνική Ορνιθολογική Εταιρεία, Αθήνα.
- Muller, J., Pollath, J., Moshammer, R. & Schroder, B. (2009). Predicting the occurrence of Middle Spotted Woodpecker *Dendrocopos medius* on a regional scale, using forest inventory data. *Forest Ecology and Management* 257: 502-509.
- Muller, M., Pasinelli, G., Schiegg, K., Spaar, R. & Jenni, L. (2005). Ecological and social effects on reproduction and local recruitment in the red-backed shrike. *Oecologia* 143: 37-50.
- Mundy, P.J., Butchart, D., Ledger, J. & Piper, S. (1992). *The Vultures of Africa*. Academic Press, London.
- Murphy, E.C. & Lehnhausen, W.A. (1998). Density and foraging ecology of woodpeckers following a stand-replacement fire. *Journal of Wildlife Management* 62: 1359-1372.

- Μακρυγιάννη, Ε., Ναζηρίδης, Θ., Παναγιωτοπούλου, Μ. & Τσουγκράκης, Γ. (2008). Αδημοσίευτα δεδομένα του προγράμματος LIFE – Nature “Conservation of *Anser erythropus* on European migration route”.
- Naccari, C., Cristani, M., Cimino, F., Arcoraci, T. & Trombetta, D. (2009). Common buzzards (*Buteo buteo*) bio-indicators of heavy metals pollution in Sicily (Italy). *Environment International* 35: 594-598.
- Navarro, J., Gonzalez-Solis, J. & Viscor, G. (2007). Nutritional and feeding ecology in Cory's shearwater *Calonectris diomedea* during breeding. *Marine Ecology Progress Series* 351: 261-271.
- Nemeckova, I., Mrlik, V. & Drozd, P. (2008), Timing of breeding, habitat preference and reproductive success of marsh harriers (*Circus aeruginosus*). *Biologia* 63: 261-265.
- Newbery, P., Schäffer, N. & Smith, K. (2001). European Union Action Plan for Bittern (*Botaurus stellaris*) European Union Action Plans for 8 Priority Birds Species.
- Newbrey, J.L., Bozek, M.A. & Niemuth, N.D. (2005). Effects of lake characteristics and human disturbance on the presence of piscivorous birds in northern Wisconsin, USA. *Waterbirds* 28: 478-486.
- Newton, I. (1979). *Population Ecology of Raptors*. Buteo Books, Vermillion.
- Nikolov, B.P. & Hristova, I.P. (2007). Time-activity budgets of juvenile Woodchat Shrikes *Lanius senator* during the post-fledging period. *Ardea* 95(2): 235-241.
- Ontiveros, D., Pleguezuelos, J.M. & Caro, J. (2005). Prey density, prey detectability and food habits: The case of Bonelli's eagle and the conservation measures. *Biological Conservation* 123: 19-25.
- Oro, D. (1995). The influence of commercial fisheries in daily activity of Audouin's gull *Larus audouinii* in the Ebro Delta, NE Spain. *Ornis Fennica* 72: 154-158.
- Oro, D., De Leon, A., Minguéz, E. & Fürness, R.W. (2005). Estimating predation on breeding European storm-petrels (*Hydrobates pelagicus*) by yellow-legged gulls (*Larus michahellis*). *Journal of Zoology* 265: 421-429, Part 4.
- Oro, D., Ruiz, X., Jover, L., Pedrocchi, V. & Gonzalez-Solis, J. (1997). Diet and adult time budgets of Audouin's Gull *Larus audouinii* in response to changes in commercial fisheries. *Ibis* 139: 631-637.
- Palomino, D. & Carrascal, L.M. (2007). Habitat associations of a raptor community in a mosaic landscape of Central Spain under urban development. *Landscape and Urban Planning* 83: 268-274.
- Panayotopoulou, M., Kokonis Y. & Chatzinikolaou Y. In: Litt. status, ecology and conservation of the White-headed Duck (*Oxyura leucocephala*) in Greece.
- Papageorgiou, N.K., Vlachos, C.G. & Bakaloudis, D. (1993). Diet and nest site characteristics of Eagle Owl (*Bubo bubo*) breeding in two different habitats in north-eastern Greece. *Avocetta* 17: 49-54.
- Pasinelli, G. (2000). Oaks (*Quercus* sp.) and only oaks? Relations between habitat structure and home range size of the middle spotted woodpecker (*Dendrocopos medius*). *Biological Conservation* 93: 227-235.
- Pasinelli, G. (2007). Nest site selection in middle and great spotted woodpeckers *Dendrocopos medius* & *D. major*: implications for forest management and conservation. *Biodiversity and Conservation* 16: 1283-1298.
- Pechacek, P. & Oleire-Oltmanns, W. (2004). Habitat use of the three-toed woodpecker in central Europe during the breeding period. *Biological Conservation* 116: 333-341.
- Pechacek, P. (2001). *Tree selection for roosting and nesting in the three-toed woodpecker: consequences for forest management*. In: Field, R, Warren, R.J., Okarma, H. & Sievert, P.R. (eds). Proceedings of the 2nd international wildlife management congress. The Wildlife Society, Bethesda. Pp 15-18.

- Pechacek, P. (2004). Spacing behavior of Eurasian three-toed woodpeckers (*Picoides tridactylus*) during the breeding season in Germany. *Auk* 121: 58-67.
- Pedrocchi, V., Oro, D., Gonzalez-Solis, J., Ruiz, X. & Jover, L. (2002). Differences in diet between the two largest breeding colonies of Audouin's gulls: The effects of fishery activities. *Scientia Marina* 66: 313-320.
- Petry, M.V., Krüger, L., Da Silva Fonseca, V.S., Brummelhaus, J., & Da Cruz Piuco, R. (2009). Diet and ingestion of synthetics by Cory's Shearwater *Calonectris diomedea* off southern Brazil. *J. of Ornithology* (in press).
- Pettersson, B. (1985). Relative importance of habitat area, isolation and quality for the occurrence of the middle spotted woodpecker *Dendrocopos medius* in Sweden. *Holarctic Ecology* 8: 53-58.
- Petty, S.J. (1999). Diet of tawny owls (*Strix aluco*) in relation to field vole (*Microtus agrestis*) abundance in a conifer forest in northern England. *Journal of Zoology* 248: 451-465.
- Pieper, H. (1981). Zur säugertiernahrung des Uhus (*Bubo bubo*) auf der griechischen Insel Lesbos. *Vogelwelt* 102: 55-56.
- Poirazidis, K., Goutner, V., Skartsi, T. & Stamou, G. (2004). Modelling nesting habitat as a conservation tool for the Eurasian black vulture (*Aegypius monachus*) in Dadia nature reserve, north-eastern Greece. *Biological Conservation* 118: 235-248.
- Poirazidis, K., Goutner, V., Tsachalidis, E. & Kati, V. (2007). Comparison of nest site selection patterns of different sympatric raptor species as a tool for their conservation. *Animal Biodiversity and Conservation* 30: 131-145.
- Pons, P. & Bas, J.M. (2005). Open-habitat birds in recently burned areas: the role of the fire extent and species habitat breadth. *Ardeola* 52: 119-131.
- Pons, P. & Prodon, R. (1996). Short term temporal patterns in a Mediterranean shrubland bird community after wildfire. *Acta Oecologica* 17: 29-41.
- Pons, P., Lambert, B., Rigolot, E. & Prodon, R. (2003). The effects of grassland management using fire on habitat occupancy and conservation of birds at a mosaic landscape. *Biodiversity and Conservation* 12: 1843-1860.
- Poulsen, J.G. (1996). Behaviour and parental care of Skylark *Alauda arvensis* chicks. *Ibis* 138: 525-531.
- Poulsen, J.G., Sotherton, N.W. & Aebischer, N.J. (1998). Comparative nesting and feeding ecology of skylarks *Alauda arvensis* on arable farmland in southern England with special reference to set-aside. *Journal of Applied Ecology* 35: 131-147.
- Prieto, J., Gonzalez-Solis, J., Ruiz & Jover X.L. (2003). Can rats prey on gull eggs? An experimental approach. *Biodiversity and Conservation* 12: 2477-2486.
- Radović, A. & Mikuska T. (2009). Population size, distribution and habitat selection of the white-tailed eagle *Haliaeetus albicilla* in the alluvial wetlands of Croatia. *Biologia* 64: 156-164.
- Ratcliffe, D. (1993). *The peregrin falcon*. T. & A.D. Poyser, London.
- Redpath, S., Madders, M., Donnelly, E., Anderson, B., Thirgood, S., Martin, A. & Mcleod M. (1998). Nest site selection by Hen Harriers in Scotland. *Bird Study* 45: 51-61.
- Ristow, D. & Wink, M. (1985). *Breeding success and conservation management of Eleonora's Falcon*. In: Newton, I. & Chancellor, R. (eds.). Conservation Studies of Raptors. ICBP Technical Publication No 5, Cambridge, UK. Pp. 147-152.
- Ristow, D. & Xirouchakis, S. (2000). What is killing Eleonora's Falcons? *World Bird Watch* 22: 14-15.
- Ristow, D. (2001). Poison is causing the sudden population decline of Eleonora's Falcon. *Intern Hawkwatcher* 3: 10-17.

- Ristow, D., Conrad, B., Wink, C. & Wink, M. (1980). Pesticide residues of failed eggs of Eleonora's Falcon (*Falco eleonora*) from Aegean colony. *Ibis* 122: 74-76.
- Ristow, D., Feldmann, F., Scharlau, W., Wink, C., & Wink, M. (1991). *Population dynamics of Cory's Shearwater (Calonectris diomedea) and Eleonora's falcon (Falco eleonora) in Eastern Mediterranean*. In: Seitz, A., Loeschcke, V. (eds). *Species conservation: A Population-Biological Approach*. Birkhauser Verlag, Basel. Pp 199-212.
- Ristow, D., Wink C., Wink, M. (1986). Assessment of Mediterranean Autumn Migration by Prey Analysis of Eleonora's Falcon. Proc. 1st Conf. on Birds wintering in the Mediterranean Region, Aulla Feb. 1984. *Supplemento alle Ricerche di Biologia della Selvaggina* 10(1): 285-295.
- Rizzolli, F., Sergio, F., Marchesi, L. & Pedrini, P. (2005). Density, productivity, diet and population status of the Peregrine Falcon *Falco peregrinus* in the Italian Alps. *Bird Study* 52: 188-192.
- Roberge, J.M., Angelstam, P., & Villard, M.A. (2008). Specialised woodpeckers and naturalness in hemiboreal forests – Deriving quantitative targets for conservation planning. *Biological conservation* 141: 997-1012.
- Robles, H., Ciudad, C., Vera, R. & Baglione, V. (2007). No effect of habitat fragmentation on post-fledging, first-year and adult survival in the middle spotted woodpecker. *Ecography* 30: 685-694.
- Robles, H., Ciudad, C., Vera, R., Olea, P.P., Purroy, F.J. & Matthysen, E. (2007). Sylvopastoral management and conservation of the middle spotted woodpecker at the south-western edge of its distribution range. *Forest Ecology and Management* 242: 343-352.
- Rosenvald, R. & Löhmus, A. (2003). Nesting of the black stork (*Ciconia nigra*) and white-tailed eagle (*Haliaeetus albicilla*) in relation to forest management. *Forest Ecology and Management* 185: 217-223.
- Ruben, F., Krijgsveld, K., Heunks C., Poot, M. & Dirksen, S. (2009). *Nocturnal and Diurnal Flight Intensity and Altitude of Seabirds and Migrants in and around an Offshore WindFarm in the Dutch North Sea in Zeebrugge, Belgium*. In: Stienen, E., Ratcliffe, N., Seys, J., Tack, J., Mees, J. & Dobbelaere, I. (eds.). 2009. Seabird Group 10th International Conference. Provincial Court, Brugge, 27-30 March 2009. VLIZ Special Publication 42. Communications of the Research Institute for Nature and Forest-INBO.M.2009.1. Research Institute for Nature and Forest (INBO), Brussels, Belgium-Flanders Marine institute (VLIZ). Oostende, Belgium. Viii+68 p.
- Rudolini, D., Gustin, M., Bogliani, G. & Garavaglia, R. (2005). Birds and powerlines in Italy: an assessment. *Bird Conservation International* 15(2): 131-145.
- Ruiz, C., Schindler, S. & Poirazidis, K. (2005). *Impact of Wind Farms on Birds in Thrace, Greece*. Technical Report, WWF-Greece, Athens.
- Sackl, P., Tiefenbach, M., Ilzer, W., Pfeiler, J. & Wieser, B. (2004). Monitoring the Austrian relict population of European roller *Coracias garrulus* - a review of preliminary data and conservation implications. *Acrocephalus (Ljubljana)* 121: 51-57.
- Sanchez, S., Václav, R. & Prokop, P. (2009). An inter-regional approach to intraspecific variation in habitat association: Rock Buntings *Emberiza cia* as a case study. *Ibis* 151: 88-98.
- Sánchez-Zapata, J.A., Carrete, M., Gravilov, A., Sklyarenko, S., Ceballos, O., Donázar, J.A. & Hiraldo, F. (2003). Land use changes and raptor conservation in steppe habitats of Eastern Kazakhstan. *Biological Conservation* 111: 71-77.
- Sanders, M.D. & Maloney, R.F. (2002). Causes of mortality at nests of ground-nesting birds in the Upper Waitaki Basin, South Island, New Zealand: A 5-year video study. *Biological Conservation* 106: 225-236.
- Sanz, A., Minguez, E., Anadon, J.D. & Hernandez, V.J. (2005). Heterogeneous use of space in three breeding territories of bonelli's eagle (*Hieraetus fasciatus*). *Ardeola* 52: 347-350.

- Sato, K., Daunt, F., Watanuki, Y., Takahashi, A. & Wanless, S. (2008). A new method to quantify prey acquisition in diving seabirds using wing stroke frequency. *J. of Exp. Biology* 211: 58-65.
- Saurola, P.L. (1997). The osprey (*Pandion haliaetus*) and modern forestry: A review of population trends and their causes in Europe. *Journal of Raptor Research* 31: 129-137.
- Schwemmer, P. & Stefan, Z. (2006). Spatial patterns in at-sea behaviour during spring migration by little gulls (*Larus minutus*) in the southeastern North Sea. *Journal of Ornithology* 147(2): 354-366.
- Scozzafava, S. & De Sanctis, A. (2006). Exploring the effects of land abandonment on habitat structures and on habitat suitability for three passerine species in a highland area of Central Italy. *Landscape and Urban Planning* 75: 23-33.
- Selås, V., Tveiten, R. & Aanonsen, O.M. (2007). Diet of Common Buzzards (*Buteo buteo*) in southern Norway determined from prey remains and video recordings. *Ornis Fennica* 84: 97-104.
- Selva, N., Jerzejewska, B., Jedrzejewski, W. & Wajrak, A. (2005). Factors affecting carcass use by a guild of scavengers in European temperate woodland. *Canadian Journal of Zoology* 83: 1590-1601.
- Sergio, F. & Boto, A. (1999). Nest dispersion, diet, and breeding success of Black Kites (*Milvus migrans*) in the Italian pre-Alps. *Journal of Raptor Research* 33: 207-217.
- Sergio, F., Boto, A., Scandolara, C. & Bogliani, G. (2002). Density, nest sites, diet, and productivity of Common Buzzards (*Buteo buteo*) in the Italian pre-Alps. *Journal of Raptor Research* 36: 24-32.
- Sergio, F., Scandolara, C., Marchesi, L., Pedrini, P. & Penteriani, V. (2005). Effect of agro-forestry and landscape changes on common buzzards (*Buteo buteo*) in the Alps: Implications for conservation. *Animal Conservation* 8: 17-25.
- Serrano, D. & Astrain, C. (2005). Microhabitat use and segregation of two sibling species of Calandrella larks during the breeding season: Conservation and management strategies. *Biological Conservation* 125: 391-397.
- Shamoun-Baranes, J., Van Loon, E., Alon, D., Alpert, P., Yom-Tov, Y. & Leshem, Y. (2006). Is there a connection between weather at departure sites, onset of migration and timing of soaring-bird autumn migration in Israel? *Global Ecology and Biogeography* 15: 541-552.
- Shealer, D.A., Buzzell, J.M. & Hejar, J.P. (2006). Effect of floating nest platforms on the breeding performance of Black Terns *Journal of Field Ornithology* 77(2).
- Shrubb, M. (1997). Historical trends in British and Irish corn buntings *Miliaria calandra* populations – evidence for the effects of agricultural change. In: Donald, P.F. & Aebischer, N.J. (eds). *The Ecology and Conservation of Corn Buntings Miliaria Calandra*. Joint Nature Conservation Committee, Peterborough, UK. Pp. 27-41.
- Sidorovich, V.E., Solovej, I.A., Sidorovich, A.A. & Rotenko, I. (2008). Effect of felling on the distribution of rodents and their predators in a transitional mixed forest. *Polish Journal of Ecology* 56: 309-321.
- Sielicki, J. & Mizera, T. (eds). (2009). *Peregrine Falcon populations - status and perspectives in the 21st century*. European Peregrin Falcon Group - Society for the Protection of Wild Animals "Falcon". Turul, Warsaw.
- Siokhin, V. (1993). Factors influencing the population structure and trophic levels in the main breeding colonies of gulls and terns in the Black and Azov seas. In: Aguilar, J.S., Monbailliu, X. & Paterson, A.M. *Status and Conservation of Seabirds*. Proceedings of the 2nd Medmaravis. SEO, Madrid.
- Sirami, C., Brotons, L. & Martin, J.L. (2007). Vegetation and songbird response to land abandonment: from landscape to census plot. *Diversity and Distributions* 13: 42-52.

- Siriwardena, G.M., Baillie, S.R., Crick, H.Q.P. & Wilson, J.D. (1999). The importance of variation in the breeding performance of seed-eating birds in determining their population trends on farmland. *Journal of Applied Ecology* 37: 1-22.
- Skartsi, T., Elorriaga, J., Vasilakis, D. & Poirazidis, C. (2008). Population, breeding and conservation status of Eurasian Black Vulture in the Dadia National Park, Thrace, NE Greece. *Journal of Natural History* 42: 345-353.
- Skartsi, T., Vasilakis, D. & Elorriaga, J. (2008). *Population trends and conservation of vultures in the National Park of Dadia-Lefkimi-Soufli forest*. In: Catsadorakis, G. (ed.) The Dadia - Lefkimi - Soufli National Park, Greece: Biodiversity, Management and Conservation. WWF-Hellas, Athens. (in press).
- Slotta-Bachmayr, L. Bögel, R. & Camina, C.A. (eds). (2004). *The situation of Griffon Vulture in Greece*. In: The Eurasian Griffon Vulture (*Gyps fulvus*) in Europe and the Mediterranean. Status report and Action plan.
- Solonen, T. & Ursin, K.A. (2008). Breeding of Tawny Owls *Strix aluco* in rural and urban habitats in southern Finland. *Bird Study* 55: 216-221.
- Steiner, H. (2000). Forest fragmentation, competition and climatic dependence in the Honey Buzzard (*Pernis apivorus*). *Journal für Ornithologie* 141: 68-76.
- Stevens, D.K., Donald, P.F., Evans, A.D., Buckingham, D.L. & Evans, J. (2002). Territory distribution and foraging patterns of ciril buntings (*Emberiza cirius*) breeding in the UK. *Biological Conservation* 107: 307-313.
- Stoate, C., Borralho, R. & Araújo, M. (2000). Factors affecting corn bunting *Miliaria calandra* abundance in a Portuguese agricultural landscape. *Agriculture, Ecosystems and Environment* 77: 219-226.
- Suarez, F., Yanes, M., Herranz, J. & Manrique, J. (1993). Nature reserves and the conservation of Iberian shrub-steppe passerines: The paradox of nest predation. *Biological Conservation* 64: 77-81.
- Suarez, S., Balbontín, J. & Ferrer, M. (2000). Nesting habitat selection by booted eagles *Hieraaetus pennatus* and implications for management. *Journal of Applied Ecology* 37: 215-223.
- Sulkava, S., Tornber, G.R. & Koivusaari, J. (1997). Diet of the white-tailed eagle *Haliaeetus albicilla* in Finland. *Ornis Fennica* 74: 65-78.
- Sultana, J. & Borg, J.J. (2005). *Population ecology and conservation of the European Storm-Petrel *Hydrobates pelagicus* in the Mediterranean*. In: Proceedings of the 1st Symposium on the Mediterranean Action Plan for the Conservation of Marine and Coastal Birds. Spain, 17-19 November.
- Sunde, P. & Bølstad, M.S. (2004). A telemetry study of the social organization of a tawny owl (*Strix aluco*) population. *Journal of Zoology* 263: 65-76.
- Svehlik, J., Meyburg, B-U. (1979). Clutch size and breeding success of the Lesser Spotted Eagle (*Aquila pomarina*) and the Imperial Eagle (*Aquila beliaca*) in the Carpathians of Eastern Slovakia 1966-1978. *Journal of Ornithology* 120: 406-415.
- Taylor, E.C., Green, R.E. & Perrins, J. (2007). Stone-curlews *Burhinus oedipnemus* and recreational disturbance: developing a management tool for access. *Ibis* 149(1): 37-44.
- Tella, J.L. & Forero, M.G. (2000). Farmland habitat selection of wintering lesser kestrels in a Spanish pseudosteppe: Implications for conservation strategies. *Biodiversity and Conservation* 9: 433-441.
- Thalmann, S., Baker, G.B., Hindell, M., Double, M.C. & Gales, R. (2007). Using biometric measurements to determine gender of Flesh-footed Shearwaters, and their application as a tool in long-line by-catch management and ecological field studies. *Emu* 107: 231-238.
- Thibault, J.C., Vigne, J.D. & Torre, J. (1992). The diet of young Lammergeiers *Gypaetus barbatus* in Corsica: its dependence on extensive grazing. *Ibis* 135: 42-48.

- Thirion, J.-M. & Lebon, P. (2006). Territory and daily rhythms of male Tawny pipit *Anthus campestris* from the Charente-Maritime coastal population. *Alauda* 74(3): 323-330.
- Thompson, S., Hazel, A., Bailey, N., Bayliss, J. & Lee, J.T. (2004). Identifying potential breeding sites for the stone curlew (*Burhinus oedipnemos*) in the UK. *Journal for Nature Conservation* 12, 229 - 235.
- Tores, M. & Y. Yom-Tov (2003) The diet of the barn Owl *Tyto Alba* in the Negev desert. *Israel Journal of Zoology* 49: 233-236.
- Toschik, P.C., Christman, M.C., Rattner, B.A. & Ottinger, M.A. (2006). Evaluation of osprey habitat suitability and interaction with contaminant exposure. *Journal of Wildlife Management* 70: 977-988.
- Trierweiler, C., Drent, R.H., Komdeur, J., Exo, K.-M., Bairlein, F. & Koks, B.J. (2008). The annual cycle of Montagu's Harrier *Circus pygargus*: Driven by voles and grasshoppers. *Limosa* 81: 107-115.
- Tsatsakis, A., Christakis-Hampsas, M., Xirouchakis, S., Baum, F. & Ristow, D. (2001). Whodunit? The case of the disappearing Eleonora's Falcons. *World Bird Watch* 23: 25-27.
- Tucker, G. M. & Heath, M. F. (1994). *Birds in Europe: Their conservation status*. BirdLife International (BirdLife Conservation Series No 3), Cambridge, UK.
- Ursúa, E., Serrano, D. & Tella, J.L. (2005). Does land irrigation actually reduce foraging habitat for breeding lesser kestrels? The role of crop types. *Biological Conservation* 122: 643-648.
- Vagliano (1981). *Contribution au statut des rapaces diurnes et nocturnes nicheurs en Crete*. In: *Rapaces Méditerranéées*, Annales du CROP, 1. Aix en Provence. Pp. 14-16.
- Valera, F., Kristin, A. & Hoi, H. (2001). Why does the lesser grey shrike (*Lanius minor*) seldom store food? Determinants of impaling in an uncommon storing species. *Behaviour* 138(11-12): 1421-1436.
- Van Turnhout, C. (2005). The disappearance of the Tawny Pipit *Anthus campestris* as a breeding bird from the Netherlands and Northwest-Europe. *Limosa* 78(1): 1-14.
- Vangeluwe, D., Handrinos, G. & Bulteau, V. (1998). Le point sur le Courlis à bec grele (*Numenius tenuirostris*). Ou l'observer, comment l'identifier ? *Ornithos* 5: 22-35.
- Vanhinsbergh, D. & Evans, A. (2002). Habitat associations of the Red-backed Shrike (*Lanius collurio*) in Carinthia, Austria. *J. Ornithol.* 143: 405-415.
- Vepsäläinen, V., Pakkala, T., Piha, M. & Tiainen, J. (2005). Population crash of the Ortolan Bunting *Emberiza hortulana* in agricultural landscapes of southern Finland. *Ann. Zool. Fennici* 42: 91-107.
- Village, A. (1990). *The Kestrel*. A. & A.D. Poyser, London.
- Vlachos, C., Bakaloudis, D. & Chatzinikos, E. (2004). Unusual nesting of the Lesser Kestrel (*Falco naumanni*) in Thessaly, Greece. *J. Rap. Research* 38: 161-163.
- Vlachos, C., Bakaloudis, D., Chatzinikos, E., Papadopoulos, T. & Tsalagas D. (2003). Aerial hunting behaviour of the Lesser Kestrel *Falco naumanni* during the breeding season in Thessaly (Greece). *Acta Ornithologica* 38: 129-134.
- Voskamp, P. (2000). Population biology and landscape use of the Honey Buzzard *Pernis apivorus* in Salland. *Limosa* 73: 67-76.
- Wakeham-Dawson, A., Szoszkiewicz, K., Stern, K. & Aebischer, N.J. (1998). Breeding skylarks *Alauda arvensis* on environmentally sensitive area arable reversion grass in southern England: survey-based and experimental determination of density. *Journal of Applied Ecology* 35: 635-648.
- Walter, H. (1979). *Eleonora's Falcon: adaptations to prey and habitat in a social raptor*. University of Chicago Press, Chicago and London.

- Warham, J. (1990). *The Petrels: their ecology and breeding systems*. Academic Press, London.
- Watanuki, Y., Daunt, F., Takahashi, A., Newell, M., Wanless, S., Sato, K. & Miyazaki, N. (2008). Microhabitat use and prey capture of a bottom-feeding top predator, the European shag, shown by camera loggers. *Marine Ecology Progress Series* 356: 283-293.
- Watson J., Leitch, A.F. & Rae, S.R. (1993). The diet of the Golden Eagle *Aquila chrysaetos* in Scotland. *Ibis* 135: 387-393
- Watson, J. (1997). *The Golden Eagle*. T. & A.D. Poyser, London, U.K.
- Watts, B.D. & Paxton, B.J. (2007). Ospreys of the Chesapeake Bay: Population recovery, ecological requirements, and current threats. *Waterbirds* 30 (Spec. Iss.): 39-49.
- Wetlands International. (2006). *Waterbird Population Estimates - 4th edition*. Wetlands International. Wageningen.
- Wheeler, P. (2008). Effects of sheep grazing on abundance and predators of field vole (*Microtus agrestis*) in upland Britain. *Agriculture, Ecosystems and Environment* 123(1-3): 49-55.
- White G., Purps J. & Alsbury, S. (2006). *The Bittern in Europe: a guide to species and habitat management*. The RSPB.
- Wiacek, J. (2008). Benefits and costs of semi-colonial breeding in the Montagu's Harrier *Circus pygargus*. *Belgian Journal of Zoology* 138: 36-40.
- Wiemeyer, S.N., Lamont, T.G. & Locke, L.N. (1980). Residues of environmental pollutants and necropsy data for Eastern United States Ospreys, 1964-1973. *Estuaries* 3: 155-167.
- Wilson, J.D., Evans, J., Browne, S.J. & King, J.R. (1997). Territory distribution and breeding success of skylarks *Alauda arvensis* on organic and intensive farmland in southern England. *Journal of Applied Ecology* 34: 1462-1478.
- Wink, M. & Ristow, D. (2000). *Biology and molecular genetics of Eleonora's falcon Falco eleonora, a colonial raptor of Mediterranean islands*. In: Chancellor, R.D., Meyburg, B.-U. (eds). *Raptors at Risk*. World Working Group on Birds of Prey/ Hancock House, Surrey. Pp 653-668.
- Wirtitsch, M., Hoi, H., Valera, F. & Kristin, A. (2001). Habitat composition and use in the lesser grey shrike (*Lanius minor*). *Folia Zoologica* 50(2): 137-150.
- Wuczynski, A. (2005). Habitat use and hunting behaviour of Common Buzzards *Buteo buteo* wintering in south-western Poland. *Acta Ornithologica* 40: 147-154.
- Xirouchakis, S. & Tsiakiris, R. (2009). *Status and population trends of vultures in Greece*. In: Donazar, J.A. & Margalida, A. (eds.). *Vulture conservation and carcasses management*. (in press).
- Xirouchakis, S. (2001). The Golden eagle *Aquila chrysaetos* in Crete. Distribution, population status and conservation problems. *Avocetta* 25: 275-281.
- Xirouchakis, S. (2003). *Population trends and aspects of breeding biology of the Bearded Vulture Gypaetus barbatus in Crete (1996-2002)*. Pp 61-67. In: Sarrazin, F. & Thiollay, J.-M. (Eds.). *Proceedings of the international meeting. Conservation and management of Bearded Vulture populations*. Ligue pour la protection des Oiseaux (LIPO). Tende, France.
- Xirouchakis, S. (2004). *Causes of raptor mortality in Crete*. Pp: 849-860. In: Meyburg, B.-U. & Chancellor, R. (Eds). *Raptors Worldwide*. WWGBP/ MME.
- Xirouchakis, S. (2005). The avifauna of the western Rodopi forests (N. Greece). *Belgian Journal of Zoology* 135: 261-269.
- Xirouchakis, S., Grivas, C., Probonas, M., Sakoulis, A. & Andreou, G. (2003). *Evaluation of actions for the conservation of Bearded Vulture (Gypaetus barbatus) in Crete*. Pp. 124-

132. In: Sarrazin, J-F. & Thiollay, J-M. (Eds). Proceedings of the international meeting. Conservation and management of Bearded Vulture populations. Ligue pour la protection des Oiseaux (LIPO). Tende, France,
- Xirouchakis, S., Sakoulis, A. & Andreou, G. (2001). The decline of the Bearded vulture *Gypaetus barbatus* in Greece. *Ardeola* 48: 183-190.
- Xirouchakis, S.M & Tsiakiris, R. (2009). *Status and population trends of vultures in Greece*. In: Donazar, J.A. & Margalida, A. (Eds): Vulture conservation and carcass management. (In press). Cramp & Simmons.
- Xirouchakis, S.M. & Andreou, G. (2009). Foraging behaviour and flight characteristics of griffon vultures (*Gyps fulvus*) in the island of Crete (Greece). *Wildlife Biology* 15: 1-16.
- Xirouchakis, S.M. & Mylonas, M. (2004). Griffon Vulture (*Gyps fulvus*) distribution and density in Crete. *Israel Journal of Zoology* 50: 341-354.
- Xirouchakis, S.M. & Mylonas, M. (2005). Status and structure of the Griffon Vulture (*Gyps fulvus*) population in Crete. *European Journal of Wildlife Research* 51: 223-231.
- Xirouchakis, S.M. (2005). The diet of the Griffon Vulture (*Gyps fulvus*) in Crete. *Journal of Raptor Research* 39: 179-181.
- Χανδρινός, Γ. (1992). Πουλιά. Στο: Καρανδεινός, Μ. & Λεγάκις, Α. Το Κόκκινο Βιβλίο των απειλούμενων Σπονδυλόζων της Ελλάδας. Ελληνική Ζωολογική Εταιρεία & Ελληνική Ορνιθολογική Εταιρεία.
- Yanes, M., Herranz, J. & Suarez, F. (1996). Nest microhabitat selection in larks from a European semi-arid shrub-steppe: the role of sunlight and predation. *Journal of Arid Environments* 32: 469-478.
- Zawadzka, D. (1999). Feeding habits of the Black Kite *Milvus migrans*, Red Kite *Milvus milvus*, White-tailed Eagle *Haliaeetus albicilla* and Lesser Spotted Eagle *Aquila pomarina* in Wigry National Park (NE Poland). *Acta Ornithologica* 34: 65-75.
- Zogaris, S. & Handrinos, G. (2002). *The breeding status of the Ferruginous Duck (Aythya nyroca) in Greece and habitat use at its national stronghold*. In: Petkov, N., Hughes B. & Gallo-Orsi U. (eds). Ferruginous Duck: from research to conservation. Proceed. Intern. Meeting. Sofia, Bulgaria. BSPB Conservation, Series no 6. Pp 67-71.
- Ζόγκαρης, Σ., Παπανδρόπουλος, Δ., Αλιβιζάτος, Χ., Ρήγας, Ι., Χατζηρβασάνης, Β. & Καρδακάρη, Ν. (2003). Απειλούμενα Πουλιά στον Αμβρακικό. ΟΙΚΟΣ ΕΠΕ, ETANAM ΑΕ.